



National Chemical Contaminants Programme

Dairy Milk Powder
Radionuclide Results (2013/14 and 2014/15)

MPI Technical – Paper No: 2017/24

By MPI Regulation and Assurance Branch

ISBN No: 978-1-77665-536-6 (online)

ISSN No: 2253-3923 (online)

March 2017

Disclaimer

While every effort has been made to ensure the information in this publication is accurate, the Ministry for Primary Industries does not accept any responsibility or liability for error of fact, omission, interpretation or opinion that may be present, nor for the consequences of any decisions based on this information.

Requests for further copies should be directed to:

Publications Logistics Officer
Ministry for Primary Industries
PO Box 2526
WELLINGTON 6140

Email: brand@mpi.govt.nz

Telephone: 0800 00 83 33

Facsimile: 04-894 0300

This publication is also available on the Ministry for Primary Industries website at <http://www.mpi.govt.nz/news-and-resources/publications/>

© Crown Copyright - Ministry for Primary Industries

Contents

Page

| | | |
|----------|-------------------------------------------|----------|
| 1 | Summary | 1 |
| 2 | What We Tested | 1 |
| 3 | What We Found | 2 |
| 3.1 | Radionuclide activity | 2 |
| 3.2 | Interpretation of results | 3 |
| 4 | Conclusion | 4 |
| 5 | Appendices | 5 |
| 5.1 | Results for detections by year and region | 5 |

1 Summary

This National Chemical Contaminants Programme report provides results for radionuclides in a range of milk powders, sampled over the 2013/14 and 2014/15 dairy seasons. This report will be updated with results for subsequent years, as those results become available.

New Zealand's lack of nuclear facilities, and geographical isolation in the south of the Pacific Ocean, results in it being one of the countries that is least affected by anthropogenic radionuclide contamination. Nevertheless, to satisfy international certification requirements New Zealand milk powders are routinely monitored for anthropogenic radionuclide contamination in conjunction with the national survey undertaken by the National Centre for Radiation Science. Routine monitoring includes the nuclear fission product Caesium-137 as well as the transuranics Plutonium-239/240 and Americium-241. Monitoring may also include Iodine-131, Caesium-134, and Strontium-90. Additional radionuclide testing may be undertaken on occasion to meet the requirements of particular markets.

The monitoring results continue to confirm the suitability of existing production and processing practices and the suitability of the New Zealand environment with respect to ensuring New Zealand dairy products meet international radionuclide standards.

2 What We Tested

65 dairy milk powder samples and 7 liquid milk samples were tested. Liquid milk samples were tested for the presence of gamma isotopes (I-131, Cs-134, Cs-137) when milk powder availability is low during June and July. These samples were collected from manufacturers in three New Zealand regions (Waikato, Taranaki and Westland) over the 2013/14 and 2014/15 dairy seasons.

Results for I-131, Cs-134, Cs-137, Pu-239/240 and Am-241 were reported under the testing laboratory's scope of accreditation. Sr-90 results were also reported.

3 What We Found

3.1 RADIONUCLIDE ACTIVITY

Of the 366 individual radioactivity content test results, all radionuclides were reported below the calculated minimum detectable activity limits except for Cs-137 which was the only detectable artificial radionuclide. The radionuclides analysed that were below the minimum detectable activity are presented in Table 1.

Table 1: Radionuclides Analysed that were below Minimum Detectable Activity in Milk Powder for 2013/14 and 2014/15

| Radionuclide | Decay type | Half-life [#] | Analysis method | Average MDA (Bq/kg) |
|------------------------|------------|-------------------------|-------------------------------|---------------------|
| Strontium-90 (Sr-90) | β- | 28.8 y | Liquid scintillation counting | 1.69 |
| Iodine-131 (I-131) | β- | 8.0 d | γ spectroscopy | 0.31 |
| Caesium-134 (Cs-134) | β- | 2.1 y | γ spectroscopy | 0.38 |
| Plutonium-239 (Pu-239) | α | 2.4 x 10 ⁴ y | α spectrometry | 0.07* |
| Plutonium-240 (Pu-240) | α | 6563 y | α spectrometry | 0.07* |
| Americium-241 (Am-241) | α | 432 y | α spectrometry | 0.07 |

Notes

d: days; y: years

* Pu-239 and Pu-240 activities are reported as a sum of both isotopes.

Table 2 provides a summary of annual average quantified Cs-137 activity concentrations in milk powder over the 2013/14 and 2014/15 dairy seasons. Appendix 1 includes all the results (detections below and above the calculated minimum detectable activity limits) by year and region for Cs-137.

Table 2: Annual Average Cs-137 Activity Concentrations in Milk Powder for 2013/14 and 2014/15

| Region | Cs-137 ± 2σ Uncertainty (Bq/kg)* | |
|-------------|----------------------------------|------------------------|
| | Annual average 2013/14 | Annual average 2014/15 |
| Waikato | 0.41 ± 0.10 | 0.45 ± 0.21 |
| Taranaki | 0.71 ± 0.20 | 0.88 ± 0.22 |
| Westland | 0.26 ± 0.13 | 0.28 ± 0.05 |
| New Zealand | 0.48 ± 0.14 | 0.61 ± 0.22 |

Note

* This is the average of samples with results above the minimum detectable activity only.

The Cs-137 detections are very close to the calculated minimum detectable activity limits (refer Appendix 1) and are far below levels that would give rise to health concerns. The results indicate that the dairy products manufactured are safe and wholesome.

3.2 INTERPRETATION OF RESULTS

Codex Alimentarius have set Guideline limits for use following a nuclear accident (Table 3).

Table 3: Codex Alimentarius Guideline Limits for Radionuclides

| Product name | Representative radionuclides | Level (Bq/kg)* |
|-------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------|----------------|
| Infant food | ²³⁸ Pu, ²³⁹ Pu, ²⁴⁰ Pu, ²⁴¹ Am | 1 |
| Infant food | ⁹⁰ Sr, ¹⁰⁶ Ru, ¹²⁹ I, ¹³¹ I, ²³⁵ U | 100 |
| Infant food | ³⁵ S, ⁶⁰ Co, ⁸⁹ Sr, ¹⁰³ Ru, ¹³⁴ Cs, ¹³⁷ Cs, ¹⁴⁴ Ce, ¹⁹² Ir | 1000 |
| Infant food | ³ H, ¹⁴ C, ⁹⁹ Tc | 1000 |
| Foods other than infant foods | ²³⁸ Pu, ²³⁹ Pu, ²⁴⁰ Pu, ²⁴¹ Am | 10 |
| Foods other than infant foods | ⁹⁰ Sr, ¹⁰⁶ Ru, ¹²⁹ I, ¹³¹ I, ²³⁵ U | 100 |
| Foods other than infant foods | ³⁵ S, ⁶⁰ Co, ⁸⁹ Sr, ¹⁰³ Ru, ¹³⁴ Cs, ¹³⁷ Cs, ¹⁴⁴ Ce, ¹⁹² Ir | 1000 |
| Foods other than infant foods | ³ H, ¹⁴ C, ⁹⁹ Tc | 10000 |

Note

* Guideline levels apply to the sum of activities from representative radionuclides, and to a food after reconstitution or preparation for consumption.


The monitoring for the range of radionuclides listed on Table 3 is applicable only when exposure may have occurred following a nuclear accident. The International Commission on Radiation Protection (ICRP) dose reference level of 1 milliSievert per year is used as a suitable health based guidance value to inform the acceptability of a source of dietary ionising radiation exposure to the general public. The Codex Alimentarius guideline levels are based on not exceeding this dose level from a food constituting 10% of the diet.

A conservative calculation can be made of the contribution the activity concentrations of Cs-137 make to the dietary ionising radiation dose. The ICRP dose conversion coefficients for Cs-137 are used to convert the ingested activity concentration, in Bq/kg, into a daily dose of Sieverts per kg of milk powder. Intake of milk is assigned to the standardised liquid milk intake of 1.5 L per day used in international modelling of veterinary drug intakes (EHC 240). Finally the result is multiplied by 365 to represent a year's consumption.

Using the maximum activity concentration reported in New Zealand milk powder (1.32 Bq/kg) and converting this to the liquid milk equivalent the resulting calculation derives a dose of 1.1 microSieverts (µSv) per year for children and 1.4 µSv per year for adults. This is 0.11-0.14% respectively of the ICRP dose reference level. This confirms that the doses of ionising radiation received through the trace activity concentrations of Cs-137 present in New Zealand dairy product are a negligible risk to health.

In the absence of a domestic nuclear industry the legacy contamination from international atmospheric nuclear testing is the highest contributor to New Zealand's anthropogenic radionuclide levels. Cs-137 has been monitored in milk since the 1950s in New Zealand and is seen as the best model for mid- to long-lived nuclear fission products released from atmospheric nuclear testing. The detections reported in 2013/14 and 2014/15 are consistent with the gradual pattern of decline of Cs-137 through both radioactive decay, and becoming unavailable for food chain uptake in soil (Figure 1). In reconstituted milk the activity concentrations detected would be approximately 10000 fold below the Codex guideline level for Cs-137.

Figure 1: Trends in the New Zealand Annual Average of Detected Sr-90 and Cs-137 Activity Concentrations in Milk Powder between 1961 and 2014.



Sr-90 was monitored by the National Centre for Radiation Science (formerly National Radiation Laboratory) until 2000 after which activity concentrations had decayed below the minimum detectable concentration. However, ongoing monitoring remains in place to meet trade obligations.

MPI and the Institute of Environmental Science and Research (ESR) have recently published¹ a study on dietary levels of radionuclides. The study concluded that, based on the established activity concentrations and ranges, the New Zealand diet contains activity concentrations of anthropogenic radionuclides far below the Codex Alimentarius guideline levels. In addition, the activity concentrations obtained for milk powder, support its continued use as a sentinel for monitoring fallout radionuclides in terrestrial agriculture.

4 Conclusion

These results confirm that a range of milk powders manufactured in New Zealand fall well within the guideline limits for radionuclides applied internationally. This indicates that New Zealand dairy products conform to both consumer expectations and international regulatory requirements.

¹ Pearson, A.J., et al., Natural and anthropogenic radionuclide activity concentrations in the New Zealand diet, *Journal of Environmental Radioactivity* (2015), <http://dx.doi.org/10.1016/j.jenvrad.2015.05.022>

5 Appendices

5.1 RESULTS FOR DETECTIONS BY YEAR AND REGION

Table 4: Detections by Year and Region

| Location: Waikato | | | | | |
|-------------------|---------------------------|---------------|---------------|---------------------------|---------------|
| Month | ¹³⁷ Cs (Bq/kg) | ± 2 s (Bq/kg) | Month | ¹³⁷ Cs (Bq/kg) | ± 2 s (Bq/kg) |
| July '13 | < 0.15 | | July '14 | < 0.19 | |
| August '13 | < 0.49 | | August '14 | < 0.52 | |
| September '13 | 0.27 | 0.11 | September '14 | < 0.47 | |
| October '13 | 0.38 | 0.12 | October '14 | 0.20 | 0.11 |
| November '13 | < 0.37 | | November '14 | < 0.60 | |
| December '13 | < 0.59 | | December '14 | 0.40 | 0.13 |
| January '14 | 0.28 | 0.12 | January '15 | 0.68 | 0.15 |
| February '14 | 0.53 | 0.15 | February '15 | 0.79 | 0.26 |
| March '14 | 0.51 | 0.19 | March '15 | 0.25 | 0.10 |
| April '14 | 0.41 | 0.13 | April '15 | < 0.57 | |
| May '14 | 0.47 | 0.18 | May '15 | 0.36 | 0.18 |
| June '14 | < 0.18 | | June '15 | < 0.21 | |

| Location: Taranaki | | | | | |
|--------------------|---------------------------|---------------|---------------|---------------------------|---------------|
| Month | ¹³⁷ Cs (Bq/kg) | ± 2 s (Bq/kg) | Month | ¹³⁷ Cs (Bq/kg) | ± 2 s (Bq/kg) |
| July '13 | < 1.17 | | July '14 | < 0.20 | |
| August '13 | 0.37 | 0.14 | August '14 | 0.30 | 0.16 |
| September '13 | 0.44 | 0.15 | September '14 | 0.47 | 0.19 |
| October '13 | 0.64 | 0.18 | October '14 | 0.69 | 0.15 |
| November '13 | 0.75 | 0.25 | November '14 | 0.87 | 0.18 |
| December '13 | 0.83 | 0.22 | December '14 | 1.32 | 0.34 |
| January '14 | 0.75 | 0.25 | January '15 | 1.03 | 0.24 |
| February '14 | 0.98 | 0.20 | February '15 | 1.17 | 0.27 |
| March '14 | 0.82 | 0.24 | March '15 | 1.05 | 0.27 |
| April '14 | 0.93 | 0.21 | April '15 | 1.15 | 0.25 |
| May '14 | 0.60 | 0.19 | May '15 | 0.74 | 0.19 |
| June '14 | < 0.23 | | June '15 | < 0.0052 | |

| Location: Westland | | | | | |
|--------------------|---------------------------|---------------|---------------|---------------------------|---------------|
| Month | ¹³⁷ Cs (Bq/kg) | ± 2 s (Bq/kg) | Month | ¹³⁷ Cs (Bq/kg) | ± 2 s (Bq/kg) |
| July '13 | < 0.37 | | July '14 | < 1.1 | |
| August '13 | < 0.42 | | August '14 | < 0.35 | |
| September '13 | 0.12 | 0.079 | September '14 | < 0.27 | |
| October '13 | 0.21 | 0.12 | October '14 | < 0.47 | |
| November '13 | 0.12 | 0.091 | November '14 | 0.21 | 0.072 |
| December '13 | 0.26 | 0.12 | December '14 | 0.36 | 0.075 |
| January '14 | 0.33 | 0.15 | January '15 | 0.29 | 0.11 |
| February '14 | 0.28 | 0.11 | February '15 | 0.30 | 0.08 |
| March '14 | 0.28 | 0.14 | March '15 | 0.27 | 0.077 |
| April '14 | 0.45 | 0.16 | April '15 | < 0.25 | |
| May '14 | < 0.16 | | May '15 | < 0.37 | |
| June '14 | < 0.40 | | June '15 | < 0.59 | |